

Estimating nitrogen uptake of individual roots in container- and field-grown plants using a ^{15}N -depletion approach

Astrid Volder^{A,B,H}, Laurel J. Anderson^{A,C,G}, David R. Smart^D, Arnold J. Bloom^E, Alan N. Lakso^F and David M. Eissenstat^A

^ADepartment of Horticulture, The Pennsylvania State University, University Park, PA 16802, USA.

^BDepartment of Horticultural Sciences, Texas A & M University, TAMU 2133, College Station, TX 77843, USA.

^CDepartment of Botany/Microbiology, Ohio Wesleyan University, Delaware, OH 43015, USA.

^DDepartment of Viticulture and Enology, University of California, Davis, CA 95616, USA.

^EDepartment of Plant Science, University of California, Davis, CA 95616, USA.

^FDepartment of Horticultural Sciences, Cornell University, New York State Agricultural Experiment Station, Geneva, NY 14456, USA.

^GPresent address: Department of Botany/Microbiology, Ohio Wesleyan University, Delaware, OH 43015, USA.

^HCorresponding author. Email: a-volder@tamu.edu

Abstract. We only have a limited understanding of the nutrient uptake physiology of individual roots as they age. Despite this shortcoming, the importance of nutrient uptake processes to our understanding of plant nutrition and nutrient cycling cannot be underestimated. In this study, we used a ^{15}N depletion method that allowed for the measurement of nitrate-N uptake rates on intact, individual, fine roots of known age. We expected that N uptake would decline rapidly as fine roots aged, regardless of the environmental conditions and species used. We compared age dependent uptake patterns of young grape cuttings with those of mature vines and with those of tomato. Although patterns of declining uptake with increasing root age were similar for all species and conditions tested, large differences in maximum N uptake rates existed between young cuttings and mature vines, and between woody and herbaceous species. Maximum rates were 10-fold higher for tomato and 3-fold higher for the grape cuttings, when compared with uptake rates of fine roots of mature vines. Coefficients of variation ranged from 43 to 122% within root age groups. The large variability in physiological characteristics of fine roots of the same age, diameter and order suggests that there is a functional diversity within fine roots that is still poorly understood.

Additional keywords: fine roots, nitrate uptake, nutrient uptake, root age, root diameter, root function.

Introduction

The physiology of plant nutrient acquisition under field conditions is important to both agriculture and ecology. Competition below ground for limited nutrients is a key factor influencing the success of species (e.g. Wilson 1988; Robinson *et al.* 1999), and the ability to acquire and assimilate nutrients is a key determinant of net primary production (Marschner *et al.* 1986). The ability of plants to acquire nutrients is determined by the amount of root absorptive surface area, soil conditions, and specific rates of root uptake. Understanding factors that influence the uptake of nutrients at the root surface requires a greater understanding of the traits of individual roots, as root physiology, anatomy and morphology can change dramatically depending on order, age and symbiotic association.

Many studies have focussed on the influx and efflux of water and nutrients along single root axes (Russell and

Sanderson 1967; Clarkson *et al.* 1968; Eshel and Waisel 1972, 1973; Ferguson and Clarkson 1975; Clarkson and Scattergood 1982; Henriksen *et al.* 1990; Siddiq *et al.* 1991; Wang *et al.* 1993; Kronzucker *et al.* 1995) or whole-root systems of young plants grown in hydroponic systems (Bloom 1985; Smart and Bloom 1988; Le Bot and Kirkby 1992; Raper *et al.* 1991; Andriolo *et al.* 1996). Studies performed along individual root axes generally showed that the active zone of nutrient uptake is the zone directly behind the root tip. This was interpreted as younger root tissue being more active in nutrient uptake than older root tissue, with the older root tissue located further back along the root axis. However, many absorptive roots, especially in woody plants, only extend 2 or 3 cm in the first days of life and then stop growing (e.g. Resendes *et al.* 2008). These short, very fine lateral roots age, but we have a limited understanding of their nutrient uptake physiology as they age (Lucash *et al.* 2007), particularly in light of the fact

the tip ceases growing (using an ordering scheme where roots with no laterals are 1st order) (Fitter 1982).

Such fine lateral roots are thought to be important to nutrient acquisition under field conditions. This is underscored by whole-root system studies which show that the most absorptive roots may represent only a small fraction of the total root system mass. For example, in a study comparing apparent inflow rates (using plant growth and tissue N content) and actual (measured) inflow rates, Robinson *et al.* (1991) concluded that no more than 10% of the root system of spring wheat was required for the plants' acquisition of nitrate from the soil. Similarly, Passioura (1980) estimated that the fraction of the wheat root system active in water uptake was no higher than 30%.

Others have described the root system in the context of branching orders (Macfall *et al.* 1991; Guo *et al.* 2004; Eissenstat and Volder 2005; Guo *et al.* 2008), with the most external, 1st and 2nd order roots mainly responsible for the acquisition of mineral nutrients and water, while the older, higher-order roots (e.g. >4th order) provide support, transport and anchoring as well as storage (Guo *et al.* 2008; Valenzuela-Estrada *et al.* 2009). However, 1st and 2nd order roots are themselves variable populations. A single plant will have a large number of 1st and 2nd order roots that vary widely in age (Bouma *et al.* 2001; Volder *et al.* 2005). The young cohort of 1st order roots (typically <25 mm in length) is considered the most important portion of the root system for N acquisition (Jensen 1962; Colmer and Bloom 1998; Taylor and Bloom 1998), and, as such, represent only a very small fraction of total root system mass or length. Often 1st order roots may only grow for a short duration (3–7 days) before elongation stops (e.g. Resendes *et al.* 2008). The ability of these 1st order roots to acquire N and phosphorus dramatically decreases with increasing age (Bouma *et al.* 2001; Volder *et al.* 2005). Volder *et al.* (2005) showed in a greenhouse study in grape (*Vitis rupestris* Scheele \times *V. riparia* Michx. cv. 3309C) that roots that were only 2 days old already had a 50% reduction in N uptake rate compared to that at birth. Bouma *et al.* (2001) showed strong age-related declines for phosphorus uptake in the field in apple (*Malus domestica* Borkh.) and orange (*Citrus aurantium* L.).

It is unclear whether results based on cuttings grown under greenhouse conditions (e.g. Volder *et al.* 2005) accurately reflect uptake patterns of mature plants under field conditions. Is the rapid decline observed in root functioning of greenhouse plants the same for roots of plants growing in the field? Moreover, do herbaceous annual plants show similar patterns as those observed in woody species? Part of the reason for limited research in the area is methodological. Many field techniques measuring N uptake rates have proved problematic for reasons not entirely clear (Lucash *et al.* 2005, 2007). In this study, we used a ^{15}N depletion method that allows for the measurement of uptake rates on intact, individual fine roots of known age to compare age-dependent effects of ^{15}N uptake in grape under greenhouse and field conditions. We also explored how N uptake in an annual herbaceous species (tomato) compares with that of a woody species (grape). We hypothesised that roots of seedling grapes in the greenhouse would exhibit higher N uptake rates

than those of mature woody plants in the field. We also hypothesised that N uptake of individual roots of a fast-growing herbaceous plant like tomato would be higher than that observed in individual roots of grape seedlings. In all experimental systems, we predicted strong declines in uptake with root age.

Materials and methods

Tomato experiments

Tomato (*Lycopersicon esculentum* L., cv. 174) cuttings were grown in 3.8-L pots filled with Sunshine potting mix (Sun Gro Horticulture Canada Ltd, Vancouver, Canada), either in growth chambers ($400\ \mu\text{mol m}^{-2}\ \text{s}^{-1}$ PFD, 70% RH, 25°C) at University of California, Davis (exp. 1), or in a glass greenhouse at Penn State University (exp. 2). 'Windows' of $\sim 20 \times 15$ cm (w \times h) were cut into the pots and covered with transparent, thin, acetate film. The windows were then covered with light impenetrable shade cloth. New root growth was traced on the windows daily, with different coloured permanent markers. Root growth in the windows was generally observed within 10 days after planting. After 3 weeks of root growth, an incision was made in the acetate, and a root of known-age was removed from the substrate, but kept attached to the plant. Care was taken to avoid roots that carried laterals or roots where there were doubts about root age. Roots were placed in 0.7-mL (exp. 1) or 0.4-mL (exp. 2) eppendorf vials that were filled with buffer (10 mM MES, 1 M CaSO_4 , 5 μM K_2HPO_4 at pH = 5.7) and 1 mM unlabelled KNO_3 . Vials were covered with parafilm and the window was covered with shade cloth. Roots were left to adjust to their new conditions for at least 5 h.

Following the period of adjustment, each root was gently washed with nitrogen-free buffer, and blotted dry. Organic specks were carefully removed using fine-pointed forceps. Intact 1–2 cm long roots with root tips were individually inserted into a new vial filled with a known volume of buffer and 1 mM (exp. 1) or 0.1 mM (exp. 2) K^{15}NO_3 . Thus, mainly the portion of the root that is active in N uptake was examined (Colmer and Bloom 1998; Taylor and Bloom 1998). The vials plus intact root tips were then gently taped to the window and loosely covered with parafilm to prevent evaporation, while allowing oxygen exchange with the air to prevent development of anaerobic conditions. The procedure was repeated with other roots in the window (~ 5 roots per plant, six plants total) and a vial with buffer without a root was also taped to the window of each plant as a control. The light-impenetrable cloth was placed back over the window. After the roots had been in the vial for 1 h, 10- μL samples were taken from the solutions and pipetted onto filter paper discs in a 96-well microplate. The filter paper discs were previously treated with 5 μmol of background N [$10\ \mu\text{L}$ 0.25 M $(\text{NH}_4)_2\text{SO}_4$] and allowed to dry. The microplate was kept covered until the samples were pipetted on the filter paper disc and then closed immediately. Samples taken from the uptake solution were pipetted onto new discs after 2, 4 and 6 h of root insertion, as well as five samples from the initial solution at $t=0$ h. Roots were harvested after 5 h by cutting them at the edge of the vial and rinsing them in distilled water. At the end of the

uptake period, the vials were closed immediately and the volume of the remaining solution was determined by weighing. Roots were scanned on a flatbed scanner with a transparency adaptor (WinRhizo, Regent Systems, Quebec City, Canada) and analysed for colour, length and diameter. After scanning, roots were placed in a forced air oven at 65°C and left to dry until stable mass. Root mass was determined on a microbalance; only roots with masses >0.1 mg were used for further analyses.

The paper discs were analysed for ^{15}N at the Stable Isotope Research facility for Environmental Research (SIRFER) at UC Davis. After correcting for background enrichment (unused filter discs with 5 $\mu\text{mol N}$) and volume changes in the vials, the decrease in ^{15}N content of the vial ($^{15}\text{N}_{t_0} - ^{15}\text{N}_{t_1}$) was calculated at each time point. The total decrease in ^{15}N per vial was then divided by the root mass and the time expired between the two measurement points to express uptake on a per mass per unit time basis.

Uptake = $(^{15}\text{N}_{t_0} - ^{15}\text{N}_{t_1}) / [(t_1 - t_0) \times \text{root mass}]$, where $^{15}\text{N}_{t_0}$ is the total amount of ^{15}N (nmol) in the vial at the start of the measurement period (t_0) and $^{15}\text{N}_{t_1}$ is the total amount of ^{15}N left in the vial at the end of the measurement period (t_1), $t_1 - t_0$ is the length of the measurement period in seconds, and root mass is the dry mass of the portion of the root that was in contact with the solution (mg).

In addition to vials with roots, control vials without roots were also placed on the windows and any changes in ^{15}N content were measured. The total uptake rate per vial with a root ($^{15}\text{N}_{t_0} - ^{15}\text{N}_{t_1}$) was corrected for any changes observed in the control vials over the same time period (for more information, see Volder *et al.* 2005). For most roots (except very old ones), the depletion of ^{15}N caused by root uptake greatly exceeded the average change in the control vials. For each root we also measured root length and we divided mass based uptake rates by the specific root length (mg^{-1}) to express uptake rates on a per length basis. Final solutions from exp. 1 were also analysed on a HPLC with UV detector for total NO_3^- content. This allowed for a comparison of uptake rates measured via HPLC and via the ^{15}N method (Fig. 1). These results show that uptake as measured via HPLC or using changes in $^{15}\text{N}/^{14}\text{N}$ ratio were closely correlated.

Concord grape experiments

Seedlings of grape (*Vitis labruscana* Bailey cv. Concord) were transplanted into 3.7-L pots filled with Sunshine mix (Sun Gro Horticulture Canada Ltd, Vancouver, CA, USA) and grown in the greenhouses of the Department of Horticulture at the Pennsylvania State University. The seedlings were planted in April 2000 and uptake measurements were conducted in June 2000. Root growth and nitrate-N uptake rates of individual intact fine roots were measured as described above.

For the field study, root observation boxes were placed in the root zones of mature Concord vines in April 2000 at Cornell University's Vineyard Laboratory in Fredonia, NY, USA. The vines were growing in soils that were a very deep (>3 m), very well drained, Chenango gravelly loam that was relatively uniform across the plot. Study plants were mature,




Fig. 1. Relationship between N uptake as measured via HPLC and the ^{15}N method on the same intact fine tomato roots (1 mM N initial solution). The dotted line represents a 1 : 1 line.

25-year-old *V. labruscana* cv. Concord grapevines with permanent cordons 1.8 m above the ground and spaced at 2.4 m between vines and 2.7 m between rows as described by Anderson *et al.* (2003). Roots were traced for 12 weeks until August 24 and 25 and measurements of N uptake were performed using the technique described above.

Statistics

Relationships between the two uptake methods and between vial N concentration and time were fitted using linear regression and slopes and intercepts were tested for significant deviations from 0 using the graphics software Sigmaplot 9.0 (Systat Software Inc., San Jose, CA, USA) and the JMP 7.01 statistical package (SAS Institute, Cary, NC, USA). Best fit curves for the relationships between root age, N uptake, and root diameter were determined using the graphics software Sigmaplot 9.0 (Systat Software Inc.) and tested for statistical significance using the JMP 7.01 statistical package (SAS Institute).

Results

The amount of ^{15}N in the vials generally declined linearly over time, demonstrating a constant rate of ^{15}N uptake over the measurement period (Fig. 2), suggesting that concentrations of oxygen and nitrate in the vial remained high enough to support nitrate uptake. When this decline was calculated as amount of ^{15}N taken up per gram dry root per second (Fig. 3), we found a rapid decline in ^{15}N uptake rate with increasing root age for tomato fine roots. Uptake rates had declined by 50% after 8 days. The rapid decline in uptake rate was significant, both when expressed on a mass or on a length basis (data not shown). We found a similar relationship between root age and N uptake rate for fine roots of greenhouse-grown Concord grape seedlings (Fig. 4, inset) and fine roots of mature Concord grape vines (Fig. 4). Maximum rates of uptake were lower for the mature grape vines than for the greenhouse-grown grape seedlings (Fig. 4). Maximum uptake rates for the tomato




Fig. 2. The influence of root incubation time on ^{15}N (% of original) remaining in the vials (0.4 mL) after correcting for volume changes using intact fine tomato roots (0.1 mM nitrate-N initial concentration; $n = 27$, \pm s.e.). Slopes of individual lines were used to estimate uptake rates.




Fig. 3. The relationship of N uptake with root age in intact tomato roots.




Fig. 4. The relationship of ^{15}N uptake with root age for field-grown and container-grown (inset) individual Concord grape fine roots. Note the difference in scales between the field- and container-grown dataset. The relationship for the field grown roots was: uptake = $1.30 \times \exp(-0.038 \times \text{root age})$, $R^2 = 0.233$, $P < 0.001$. The relationship for the container-grown roots was: uptake = $5.41 \times \exp(-0.072 \times \text{root age})$, $R^2 = 0.194$, $P < 0.050$.

fine roots were an order of magnitude higher than those of fine roots of field grown, mature, Concord grape (Figs 3, 4). Within species and root age classes a wide range of uptake rates was found and coefficients of variation ranged from 43 to 122% (Table 1).

There was no relationship between ^{15}N uptake and fine root diameter for either the container- or field-grown grape roots ($P > 0.24$, data not shown), but both mass-based and surface-area-based ^{15}N uptake rates of fine tomato roots declined with increasing diameter (Fig. 5). A possible underlying reason for the inverse relationship of ^{15}N uptake and diameter in tomato is that coarser roots might be older; however, we observed no relationship between root diameter and root age in tomato (Fig. 5B, inset), nor was there such a relationship for the 1st order grape roots (data not shown).

Table 1. Average ^{15}N uptake rates for several root age classes in this and earlier investigations

Average rates are mean \pm s.d.; CV, coefficient of variation

	<i>n</i>	Age range (days)	Average rate (nmol ^{15}N g _{dw} ⁻¹ s ⁻¹)	Range (nmol ^{15}N g _{dw} ⁻¹ s ⁻¹)	CV (%)	Citation
<i>Greenhouse-grown cuttings</i>						
<i>Vitis rupestris</i> \times <i>V. riparia</i> cv. 3309C	3	0–1	10.05 \pm 5.08	4.68–14.79	51	Volder et al. (2005)
<i>Vitis rupestris</i> \times <i>V. riparia</i> cv. 3309C	10	11–23	3.22 \pm 1.38	1.15–5.61	43	Volder et al. (2005)
<i>Vitis labruscana</i> cv. Concord	14	0–5	4.03 \pm 3.49	0.47–10.54	87	This investigation
<i>Vitis labruscana</i> cv. Concord	7	9–16	1.90 \pm 0.98	0.57–3.18	51	This investigation
<i>Mature field vine</i>						
<i>Vitis labruscana</i> cv. Concord	25	2–10	0.97 \pm 0.85	–0.15–3.25	88	This investigation
<i>Vitis labruscana</i> cv. Concord	64	11–77	0.44 \pm 0.54	–0.60–2.07	122	This investigation
<i>Greenhouse-grown tomato</i>						
<i>Lycopersicon esculentum</i>	19	0–3	10.60 \pm 9.18	1.92–37.20	87	This investigation
<i>Lycopersicon esculentum</i>	9	7–29	3.49 \pm 3.95	0.77–13.70	113	This investigation




Fig. 5. The relationship of ^{15}N uptake (A) per unit root surface area, and (B) per unit dry mass with root diameter for tomato roots. The regression line in (A) is: uptake = $182 \times \exp(-3.44 \times \text{root diameter})$, $R^2 = 0.448$, $P < 0.001$. The regression line in (B) is: uptake = $96.2 \times \exp(-6.2 \times \text{diameter})$, $R^2 = 0.624$, $P < 0.001$. The inset in (B) shows the lack of a relationship between root age and root diameter for tomato roots.

Discussion

Root age and nitrate uptake

In this study, we have shown that both root diameter and root age can have a large effect on the nitrate-N uptake capacity of individual roots. Both roots of young cuttings and mature vines of a woody species (grape, *V. labruscana* cv. Concord) and roots of an herbaceous species (tomato, *L. esculentum* cv. T5) showed the same general pattern: a rapid decline in nitrate-N uptake capacity as roots age. This same pattern also emerged for grape cuttings of *V. riparia* \times *V. rupestris* cv. 3309C in a previous greenhouse experiment (Volder *et al.* 2005). Maximum rates reported, however, were 10-fold higher for the herbaceous species and 3-fold higher for the seedlings, compared with uptake rates of fine roots of mature vines growing in the field. Thus, although the patterns were the same, large differences in maximum uptake existed between young cuttings and mature trees, and between woody and herbaceous species. These differences are likely related to plant growth rate and N demand (Poorter *et al.* 1990; Ter Steege *et al.* 1999), with faster growing plants such as

seedlings or herbaceous species having higher specific N uptake rates.

We found a large variability in rates of nitrate-N uptake within fine roots of the same order and age range (Table 1). There is a sizable portion of fine roots that remains relatively inactive even from the first days of birth. Other age related studies, such as those on P uptake and respiration (Bouma *et al.* 2001), root browning (Comas *et al.* 2000), N uptake and respiration (Volder *et al.* 2005), and fungal infection rate (Resendes *et al.* 2008) found similar variation in root function within fine roots of the same age, where a large proportion of the roots appears to be inactive.

Some roots, commonly called 'pioneer' or 'framework' roots that often become part of the woody root system infrastructure, may be less physiologically active in nutrient absorption and less prone to mycorrhizal colonisation than roots that are destined to remain 1st order roots (Wells and Eissenstat 2003). These pioneer roots often have a larger root tip, are born with a larger diameter, and extend in a more indeterminate fashion than those roots that are likely to remain only 1st or 2nd order and never develop secondary growth (Eissenstat and Achor 1999). Among the 1st order roots that are very fine and never undergo secondary growth, there may be distinct differences in function. For example, in apple, some of the very fine, 1st order roots are colonised within the first week by mycorrhizal fungi and have slightly higher growth rates than other similarly fine roots (Resendes *et al.* 2008). The latter may not be growing quite as actively for the first 3 days of life and soon (within 2 weeks) become colonised by non-mycorrhizal fungi or never become colonised. Thus, although new roots may be 1st order and of the same age, they may not all be destined to have the same function.

A large amount of variation in nutrient uptake rates by mature trees estimated *in situ* can be explained by the wide variety of methods employed (Lucash *et al.* 2007). Current methods, including the method employed in this paper, often do not account for diurnal variation in uptake (Scheurwater *et al.* 1999), and commonly lack adequate temperature control. However, even when the same method is employed for a short period, rates measured can vary widely (Gessler *et al.* 1998; Lucash *et al.* 2005). Nitrogen uptake rates have been reported to vary seasonally (Gessler *et al.* 1998), diurnally (Hansen 1980; Raper *et al.* 1991; Le Bot and Kirkby 1992; Cardenas-Navarro *et al.* 1998; Peuke and Jeschke 1998) and spatially along the root axis (Russell and Sanderson 1967; Henriksen *et al.* 1992; Colmer and Bloom 1998). Based upon our studies with grape cuttings, mature vines and an herbaceous plant, we conclude that nitrate-N uptake rates also can decline both as fine roots age and as diameter increases. This explains some of the seasonal variation that is reported in the literature, as maximum nitrate-N uptake rates of subsets of roots will be strongly influenced by the average age of the roots in that sample, even if the root diameter in the sample is controlled.

Both the production rate of new roots and median root lifespan will affect the average root age of a random sample of roots. During periods with high rates of root production, the root system will have a larger proportion of relatively

young root tips and this will be reflected in a random sample. Thus, during a period of high root production (e.g. late spring) we expect much higher uptake activities in an average root sample than during a period of reduced root growth (e.g. late summer), as was found by Lucash *et al.* (2005) for loblolly pine (*Pinus taeda*). In a subalpine area, where new root production may not start until the summer, Gessler *et al.* (1998) found higher uptake of ammonium during the summer than in the spring.

Root production and root longevity are affected by a great number of environmental factors, including species (Eissenstat and Yanai 1997; Black *et al.* 1998), season (Hendrick and Pregitzer 1993a), climatic conditions (Hendrick and Pregitzer 1993b), soil texture and structure (Hendrick and Pregitzer 1997), root depth (Wells and Eissenstat 2001; Baddeley and Watson 2005), plant age (Baddeley and Watson 2005) and root diameter and order (Wells and Eissenstat 2001, 2003; Baddeley and Watson 2005). The effect of root age does not only partially account for variation of N uptake measurements through time, but also in space. Roots usually have a longer lifespan deeper in the soil (Wells and Eissenstat 2001; Anderson *et al.* 2003; Joslin *et al.* 2006; Peek *et al.* 2006), increasing the average root age, which would suggest that measurements of nitrate-N uptake on random root samples from deeper soil layers would show reduced uptake rates compared with uptake rates measured on random samples from shallow soil layers. Bhat (1982) found a large decrease in nitrate uptake rates per unit root length for deeper growing apple roots in one of two apple cultivars studied, but not in the other. However, the roots used in the estimation of nitrate uptake rate in this study were growing in nutrient solution for the preceding 16 months, thus creating roots more typical of solution culture than of roots developed in soil under natural field conditions.

Implications for modelling N uptake

Plant-based ecosystem N uptake models would be improved by taking into account not only the total standing root length, but also the proportion of roots in the 0–10 day range when roots have the highest uptake capacity. Accurately quantifying and modelling fluxes of nutrients from plants to soil and *vice versa* remains one of the principal challenges in trying to predict impacts of environmental change on ecosystems (Norby and Jackson 2000). A combined index of new root production rate and standing root length (e.g. a weekly rate of new root length turnover) could be a better predictor of ecosystem nitrate-N uptake dynamics than standing root length alone.

Conclusion

Plants exhibit a rapid decline in nitrate uptake with root age regardless of growth conditions (greenhouse *v.* field) or whether the plant is woody or herbaceous. The magnitude of maximum uptake, however, varies with the environmental conditions, species, and growth demand of the plant. The large variability in physiological characteristics of fine roots of the same age and order suggests that there is a functional

diversity within fine roots that is poorly understood. The rapid decline in maximum nitrate uptake as roots age partially explains why measurements of nitrate uptake on random root samples are generally not consistent through time and space, as these samples are generally highly variable in mean root age.

Acknowledgements

We are indebted to David Harris (Stable Isotope Research facility for Environmental Research (SIRFER) at UC Davis) for performing the isotope analyses. This work was supported by USDA grant NRI 97–35107–4359 to DME, USDA/CSREES Special Research Grants Program 99–34360–7374 to DME and DRS and NSF IOS-08–18435 and USDA-NRI 2008–01029 to AJB.

References

- Anderson LJ, Comas LH, Lakso AN, Eissenstat DM (2003) Multiple risk factors in root survivorship: a 4 year study in Concord grape. *New Phytologist* **158**, 489–501. doi: 10.1046/j.1469-8137.2003.00757.x
- Andriolo JL, LeBot J, Gary C, Sappe G, Orlando P, Brunel B, Sarrouy C (1996) An experimental set-up to study carbon, water and nitrate uptake rates by hydroponically grown plants. *Journal of Plant Nutrition* **19**, 1441–1462. doi: 10.1080/01904169609365211
- Baddeley JA, Watson CA (2005) Influences of root diameter, tree age, soil depth and season on fine root survivorship in *Prunus avium*. *Plant and Soil* **276**, 15–22. doi: 10.1007/s11104-005-0263-6
- Bhat KKS (1982) Nutrient inflows into apple roots. 2. Nitrate uptake rates measured on intact roots of mature trees under field conditions. *Plant, Cell & Environment* **5**, 461–469.
- Black KE, Harbron CG, Franklin M, Atkinson D, Hooker JE (1998) Differences in root longevity of some tree species. *Tree Physiology* **18**, 259–264.
- Bloom AJ (1985) Wild and cultivated barleys show similar affinities for mineral nitrogen. *Oecologia* **65**, 555–557. doi: 10.1007/BF00379672
- Bouma TJ, Yanai RD, Elkin AD, Hartmond U, Flores-Alva DE, Eissenstat DM (2001) Estimating age-dependent costs and benefits of roots with contrasting life span: comparing apples and oranges. *New Phytologist* **150**, 685–695. doi: 10.1046/j.1469-8137.2001.00128.x
- Cardenas-Navarro R, Adamowicz S, Robin P (1998) Diurnal nitrate uptake in young tomato (*Lycopersicon esculentum* Mill.) plants: test of a feedback-based model. *Journal of Experimental Botany* **49**, 721–730. doi: 10.1093/jexbot/49.321.721
- Clarkson DT, Scattergood CB (1982) Growth and phosphate transport in barley and tomato plants during the development of, and recovery from, phosphate stress. *Journal of Experimental Botany* **33**, 865–875. doi: 10.1093/jxb/33.5.865
- Clarkson DT, Sanderson J, Russell RS (1968) Ion uptake and root age. *Nature* **220**, 805–806. doi: 10.1038/220805a0
- Colmer TD, Bloom AJ (1998) A comparison of NH_4^+ and NO_3^- net fluxes along roots of rice and maize. *Plant, Cell & Environment* **21**, 240–246. doi: 10.1046/j.1365-3040.1998.00261.x
- Comas LH, Eissenstat DM, Lakso AN (2000) Assessing root death and root system dynamics in a study of grape canopy pruning. *New Phytologist* **147**, 171–178. doi: 10.1046/j.1469-8137.2000.00679.x
- Eissenstat DM, Achor DS (1999) Anatomical characteristics of roots of citrus rootstocks that vary in specific root length. *New Phytologist* **141**, 309–321. doi: 10.1046/j.1469-8137.1999.00342.x
- Eissenstat DM, Volder A (2005) The efficiency of nutrient acquisition over the life of a root. In 'Nutrient acquisition by plants: ecological perspective'. pp. 185–220. (Springer-Verlag: Berlin)

- Eissenstat DM, Yanai RD (1997) The ecology of root lifespan. *Advances in Ecological Research* **27**, 1–60. doi: 10.1016/S0065-2504(08)60005-7
- Eshel A, Waisel Y (1972) Variations in sodium uptake along primary roots of corn seedlings. *Plant Physiology* **49**, 585–589. doi: 10.1104/pp.49.4.585
- Eshel A, Waisel Y (1973) Variations in uptake of sodium and rubidium along barley roots. *Physiologia Plantarum* **28**, 557–560. doi: 10.1111/j.1399-3054.1973.tb08606.x
- Ferguson IB, Clarkon DT (1975) Ion transport and endodermal suberization in roots of *Zea mays*. *New Phytologist* **75**, 69–79. doi: 10.1111/j.1469-8137.1975.tb01372.x
- Fitter AH (1982) Morphometric analysis of root systems – application of the technique and influence of soil fertility on root system development in two herbaceous species. *Plant, Cell & Environment* **5**, 313–322.
- Gessler A, Schneider S, Von Sengbusch D, Weber P, Hanemann U, Huber C, Rothe A, Kreuzer K, Rennenberg H (1998) Field and laboratory experiments on net uptake of nitrate and ammonium by the roots of spruce (*Picea abies*) and beech (*Fagus sylvatica*) trees. *New Phytologist* **138**, 275–285. doi: 10.1046/j.1469-8137.1998.00107.x
- Guo DL, Mitchell RJ, Hendricks JJ (2004) Fine root branch orders respond differentially to carbon source–sink manipulations in a longleaf pine forest. *Oecologia* **140**, 450–457. doi: 10.1007/s00442-004-1596-1
- Guo DL, Li H, Mitchell RJ, Han WX, Hendricks JJ, Fahey TJ, Hendrick RL (2008) Fine root heterogeneity by branch order: exploring the discrepancy in root turnover estimates between minirhizotron and carbon isotopic methods. *New Phytologist* **177**, 443–456.
- Hansen GK (1980) Diurnal-variation of root respiration rates and nitrate uptake as influenced by nitrogen supply. *Physiologia Plantarum* **48**, 421–427. doi: 10.1111/j.1399-3054.1980.tb03279.x
- Hendrick RL, Pregitzer KS (1993a) The dynamics of fine root length, biomass, and nitrogen-content in two northern hardwood ecosystems. *Canadian Journal of Forest Research* **23**, 2507–2520. doi: 10.1139/x93-312
- Hendrick RL, Pregitzer KS (1993b) Patterns of fine root mortality in two sugar maple forests. *Nature* **361**, 59–61. doi: 10.1038/361059a0
- Hendrick RL, Pregitzer KS (1997) The relationship between fine root demography and the soil environment in northern hardwood forests. *Ecoscience* **4**, 99–105.
- Henriksen GH, Bloom AJ, Spanswick RM (1990) Measurement of net fluxes of ammonium and nitrate at the surface of barley roots using ion-selective microelectrodes. *Plant Physiology* **93**, 271–280. doi: 10.1104/pp.93.1.271
- Henriksen GH, Raman DR, Walker LP, Spanswick RM (1992) Measurement of net fluxes of ammonium and nitrate at the surface of barley roots using ion-selective microelectrodes. 2. Patterns of uptake along the root axis and evaluation of the microelectrode flux estimation technique. *Plant Physiology* **99**, 734–747. doi: 10.1104/pp.99.2.734
- Jensen G (1962) Relationship between water and nitrate uptake in excised tomato root systems. *Physiologia Plantarum* **15**, 791–803. doi: 10.1111/j.1399-3054.1962.tb08128.x
- Joslin JD, Gaudinski JB, Torn MS, Riley WJ, Hanson PJ (2006) Fine-root turnover patterns and their relationship to root diameter and soil depth in a C-14-labeled hardwood forest. *New Phytologist* **172**, 523–535. doi: 10.1111/j.1469-8137.2006.01847.x
- Kronzucker HJ, Siddiqi MY, Glass ADM (1995) Kinetics of NO_3^- influx in spruce. *Plant Physiology* **109**, 319–326.
- Le Bot J, Kirkby EA (1992) Diurnal uptake of nitrate and potassium during the vegetative growth of tomato plants. *Journal of Plant Nutrition* **15**, 247–264. doi: 10.1080/01904169209364316
- Lucash MS, Joslin JD, Yanai RD (2005) Temporal variation in nutrient uptake capacity by intact roots of mature loblolly pine. *Plant and Soil* **272**, 253–262. doi: 10.1007/s11104-004-5296-8
- Lucash MS, Eissenstat DM, Joslin JD, McFarlane KJ, Yanai RD (2007) Estimating nutrient uptake by mature tree roots under field conditions: challenges and opportunities. *Trees – Structure and Function* **21**, 593–603.
- Macfall JS, Johnson GA, Kramer PJ (1991) Comparative water-uptake by roots of different ages in seedlings of loblolly pine (*Pinus taeda* L.). *New Phytologist* **119**, 551–560. doi: 10.1111/j.1469-8137.1991.tb01047.x
- Marschner H, Romheld V, Horst WJ, Martin P (1986) Root-induced changes in the rhizosphere – importance for the mineral nutrition of plants. *Zeitschrift für Pflanzenernährung und Bodenkunde* **149**, 441–456. doi: 10.1002/jpln.19861490408
- Norby RJ, Jackson RB (2000) Root dynamics and global change: seeking an ecosystem perspective. *New Phytologist* **147**, 3–12. doi: 10.1046/j.1469-8137.2000.00676.x
- Passioura JB (1980) The transport of water from soil to shoot in wheat seedlings. *Journal of Experimental Botany* **31**, 333–345. doi: 10.1093/jxb/31.1.333
- Peek MS, Leffler AJ, Hipps L, Ivans S, Ryel RJ, Caldwell MM (2006) Root turnover and relocation in the soil profile in response to seasonal soil water variation in a natural stand of Utah juniper (*Juniperus osteosperma*). *Tree Physiology* **26**, 1469–1476.
- Peuke AD, Jeschke WD (1998) The effects of light on induction, time courses, and kinetic patterns of net nitrate uptake in barley. *Plant, Cell & Environment* **21**, 765–774. doi: 10.1046/j.1365-3040.1998.00329.x
- Poorter H, Remkes C, Lambers H (1990) Carbon and nitrogen economy of 24 wild species differing in relative growth-rate. *Plant Physiology* **94**, 621–627. doi: 10.1104/pp.94.2.621
- Raper CD, Vessey JK, Henry LT (1991) Increase in nitrate uptake by soybean plants during interruption of the dark period with low intensity light. *Physiologia Plantarum* **81**, 183–189. doi: 10.1111/j.1399-3054.1991.tb02127.x
- Resendes M, Bryla D, Eissenstat D (2008) Early events in the life of apple roots: variation in root growth rate is linked to mycorrhizal and nonmycorrhizal fungal colonization. *Plant and Soil* **313**, 175–186. doi: 10.1007/s11104-008-9690-5
- Robinson D, Linehan DJ, Caul S (1991) What limits nitrate uptake from soil. *Plant, Cell & Environment* **14**, 77–85. doi: 10.1111/j.1365-3040.1991.tb01373.x
- Robinson D, Hodge A, Griffiths BS, Fitter AH (1999) Plant root proliferation in nitrogen-rich patches confers competitive advantage. *Proceedings of the Royal Society of London. Series B. Biological Sciences* **266**, 431–435. doi: 10.1098/rspb.1999.0656
- Russell RS, Sanderson J (1967) Nutrient uptake by different parts of intact roots of plants. *Journal of Experimental Botany* **18**, 491–508. doi: 10.1093/jxb/18.3.491
- Scheurwater I, Clarkson DT, Purves JV, Van Rijt G, Saker LR, Welschen R, Lambers H (1999) Relatively large nitrate efflux can account for the high specific respiratory costs for nitrate transport in slow-growing grass species. *Plant and Soil* **215**, 123–134. doi: 10.1023/A:1004559628401
- Siddiqi MY, Glass ADM, Ruth TJ (1991) Studies of the uptake of nitrate in barley. 3. Compartmentation of NO_3^- . *Journal of Experimental Botany* **42**, 1455–1463. doi: 10.1093/jxb/42.11.1455
- Smart DR, Bloom AJ (1988) Kinetics of ammonium and nitrate uptake among wild and cultivated tomatoes. *Oecologia* **76**, 336–340.
- Taylor AR, Bloom AJ (1998) Ammonium, nitrate, and proton fluxes along the maize root. *Plant, Cell & Environment* **21**, 1255–1263. doi: 10.1046/j.1365-3040.1998.00357.x
- Ter Steege MW, Stulen I, Wiersma PK, Posthumus F, Vaalburg W (1999) Efficiency of nitrate uptake in spinach: impact of external nitrate concentration and relative growth rate on nitrate influx and efflux. *Plant and Soil* **208**, 125–134. doi: 10.1023/A:1004585200558

- Valenzuela-Estrada LR, Richards JH, Diaz A, Eissenstat DM (2009) Patterns of nocturnal rehydration in root tissues of *Vaccinium corymbosum* L. under severe drought conditions. *Journal of Experimental Botany* **60**, 1241–1247. doi: 10.1093/jxb/ern367
- Volder A, Smart DR, Bloom AJ, Eissenstat DM (2005) Rapid decline in nitrate uptake and respiration with age in fine lateral roots of grape: implications for root efficiency and competitive effectiveness. *New Phytologist* **165**, 493–502. doi: 10.1111/j.1469-8137.2004.01222.x
- Wang MY, Siddiqi MY, Ruth TJ, Glass ADM (1993) Ammonium uptake by rice roots. 2. Kinetics of NH_4^+ - N^{15} influx across the plasmalemma. *Plant Physiology* **103**, 1259–1267.
- Wells CE, Eissenstat DM (2001) Marked differences in survivorship among apple roots of different diameters. *Ecology* **82**, 882–892.
- Wells CE, Eissenstat DM (2003) Beyond the roots of young seedlings: the influence of age and order on fine root physiology. *Journal of Plant Growth Regulation* **21**, 324–334. doi: 10.1007/s00344-003-0011-1
- Wilson JB (1988) A review of evidence on the control of shoot-root ratio, in relation to models. *Annals of Botany* **61**, 433–449.

Manuscript received 31 December 2008, accepted 6 May 2009