

This article was downloaded by: [Stauffer, Jr., J R][Pennsylvania State University]

On: 18 March 2011

Access details: Access Details: [subscription number 933509533]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Freshwater Ecology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smp/title-content=t925992003>

### Crush-resistance of soft-sediment gastropods of Lake Malaŵi: implications for prey selection by Molluscivorous fishes

B. N. Evers<sup>a</sup>; H. Madsen<sup>a</sup>; J. R. Stauffer Jr<sup>b</sup>

<sup>a</sup> Mandahl-Barth Research Centre for Biodiversity and Health in Developing Countries, DBL-Centre for Health Research and Development, Department of Veterinary Pathobiology, Faculty of Life Sciences, University of Copenhagen, Jaegersborg Allé 1 D, 2920 Charlottenlund, Denmark <sup>b</sup> School of Forest Resources, Penn State University, PA 16802, Pennsylvania, USA

Online publication date: 18 March 2011

**To cite this Article** Evers, B. N. , Madsen, H. and Stauffer Jr, J. R.(2011) 'Crush-resistance of soft-sediment gastropods of Lake Malaŵi: implications for prey selection by Molluscivorous fishes', *Journal of Freshwater Ecology*, 26: 1, 85 – 90

**To link to this Article: DOI:** 10.1080/02705060.2011.553921

**URL:** <http://dx.doi.org/10.1080/02705060.2011.553921>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Crush-resistance of soft-sediment gastropods of Lake Malaŵi: implications for prey selection by Molluscivorous fishes

B.N. Evers<sup>a</sup>, H. Madsen<sup>a</sup> and J.R. Stauffer Jr<sup>b\*</sup>

<sup>a</sup>Mandahl-Barth Research Centre for Biodiversity and Health in Developing Countries, DBL-Centre for Health Research and Development, Department of Veterinary Pathobiology, Faculty of Life Sciences, University of Copenhagen, Jaegersborg Allé 1 D, 2920 Charlottenlund, Denmark; <sup>b</sup>School of Forest Resources, Penn State University, University Park, PA 16802, Pennsylvania, USA

(Received 19 July 2010; final version received 1 September 2010)

An increase in human urinary schistosome transmission in southern Lake Malaŵi has been suggested to be associated with decreased density of molluscivorous fishes due to illegal seine-net fishing from the shore. In addition, the increased density of snails (*Melanoides* spp.) through the invasion of an Asian morph could have changed the predators' prey choice. At Chembe village, the intermediate host snail of urinary schistosomes, *Bulinus nyassanus*, constitutes <5% of the total gastropod fauna. This study was designed to compare crushing resistance of the intermediate host snails *Bulinus globosus* and *B. nyassanus* with that of *Melanoides tuberculata*, which dominates the gastropod fauna. A crush value index (CVI) as an indicator of potential prey value was expressed as the ratio of benefit (weight of snail tissue) to cost (crush resistance of snail shell). *Bulinus globosus* had the highest CVI. Using shell height as measure of snail size indicated that *B. nyassanus* had higher CVI than *M. tuberculata* within the size range of snails consumed by *Trematocranus placodon*, one of the molluscivore fishes. This may be one of the reasons that *B. nyassanus* is a preferred prey of *T. placodon*. In spite of this preference, the reduced population of *T. placodon* has not been able to control the population of *B. nyassanus* because of its apparent opportunistic feeding on large numbers of *M. tuberculata*.

**Keywords:** molluscivorous fishes; schistosomiasis; crush value index

### Introduction

Urinary schistosomiasis, caused by *Schistosoma haematobium* has long been recognized as a major public health problem in lakeshore communities of Lake Malaŵi (WHO 1993), where transmission has occurred primarily in streams and protected areas of the lake (Madsen et al. 2004). An increase in transmission at Cape Maclear, Lake Malaŵi, however, has occurred in recent years (Cetron et al. 1996), and Stauffer et al. (1997) suggested that this was due to an increase in abundance of intermediate host snails associated with declines in density of shallow-water molluscivorous cichlid fishes (e.g., *Trematocranus placodon*, *Trematocranus microstoma*, *Mylochromis sphaerodon*, and *Mylochromis anaphyrmus*). Prior to 1988,

---

\*Corresponding author. Email: vc5@psu.edu

the open waters of Lake Malaŵi were believed to be free from schistosome production/transmission, but Madsen et al. (2001) reported transmission along open shorelines in the southern part of the lake.

In Lake Malaŵi, there are two known intermediate hosts for *S. haematobium* – *Bulinus globosus* and *Bulinus nyassanus* (Madsen et al. 2001). *Bulinus globosus* is the primary host in sheltered areas of the lake as well as in pools and streams close to the lake shore and further inland (Madsen et al. 2004). *Bulinus nyassanus* is endemic to Lake Malaŵi and found along open shorelines on sediment of sand or gravel (Madsen et al. 2001). It has been suggested that reducing seine-net fishing in nearshore areas of the lake (Stauffer et al. 1997, 2006) would result in an increase in density (i.e., to pre-1988 levels) of molluscivorous fishes and possibly a decline in schistosome transmission. This outcome, however, requires that the molluscivores consume *B. nyassanus*.

*Trematocranus placodon* feeds on snails in the genera of both *Melanoides* and *Bulinus* (Evers et al. 2006). Stomach content analysis of field-collected *T. placodon*, however, showed that *B. nyassanus* was found in higher proportion in the stomachs than it was in the lake (Evers et al. 2006). In the environment, *B. nyassanus* constituted 3.5% of all snails, and *Melanoides* spp. comprised 94.8%. In the stomachs of *T. placodon*, the respective percentages were 24.6 and 72.3 (Evers et al. 2006). Although this would indicate a feeding preference of *T. placodon*, *Melanoides* spp. still dominate the stomach content *T. placodon*, probably because of this much greater environmental abundance. These results, however, may be confounded by the introduction of an exotic morph of *Melanoides tuberculata* (Genner et al. 2004). Indigenous *Melanoides* spp. have more sculptured shells than the introduced morph (i.e., more pronounced spiral grooves and presence of axial ribs; Genner et al. 2004) and this could represent an anti-predatory adaptation. The exotic morph has a rather smooth shell with only spiral lines; its shell may be less substantial than those of the indigenous *Melanoides* spp. Optimal foraging models indicate that prey are added to the diet in decreasing order of energetic benefit/cost ratio (Pyke et al. 1977; Begon et al. 1996). Our objective was to compare one aspect of prey suitability by comparing prey value as a crush value index (CVI) based on crushing resistance and organic dry weight (dry tissue mass) of *B. globosus*, *B. nyassanus*, and the introduced morph of *M. tuberculata*.

### Materials and methods

Snails were collected in the field during June–August 2003, and preserved in 70% ethanol. *Melanoides tuberculata* (the morph form) were collected in Lake Malaŵi at Chembe village. *Bulinus globosus* were collected from a nearby stream. The preserved snails were rinsed in distilled water, dried at 70°C for 24 h, and placed in desiccators. For *M. tuberculata*, shells that were not eroded at the apex were selected so as to measure the height accurately and test the strength of complete shells. Shell height was measured to the nearest 0.05 mm using calipers. We elected to use shell height to represent snail size although shells are crushed across their width, because *M. tuberculata* of a width comparable to those of the two *Bulinus* species would be too large for the fish to consume. Individual shell widths were not measured originally but were subsequently extrapolated from shell heights on the basis of measurements made of shell height and shell width of 25 specimens from the same samples as used for the crush resistance trials and covering the same size range for

each species ( $r^2$  values were 0.7499, 0.9216, and 0.9644 for *B. globosus*, *B. nyassanus*, and *M. tuberculata*, respectively).

Crush resistance was determined (*sensu* Brodersen et al. 2003) by placing dried snails with the aperture facing down in a Petri dish underneath a plexiglass cylinder closed at the bottom; this procedure ensured that force was applied to snails against their minimal dimension, consistent with the way the molluscivore labrid fish manipulate snails between pharyngeal plates (Stein et al. 1984). The tube was gradually filled with sand until the shell was crushed. The weight of the plexi-glass tube including sand was the crush weight. The snail and shell fragments were brushed into crucibles, and the total dry weight was determined to the nearest 0.1 mg. Crucibles containing all snail parts were combusted at 500°C for 1.5 h, and subsequently reweighed to yield inorganic weight. Organic weight (ash-free dry weight) was determined by subtracting inorganic weight from the total dry weight.

The CVI was calculated as the organic content in mg divided by crush weight in kg. Differences among species were tested using analysis of covariance ( $p < 0.05$ ) with shell height as covariate. Both the dependent variables and the covariate were  $\log_{10}$  transformed.

## Results and discussion

The characteristics of the three snail species are given in Table 1. The  $\log_{10}$  transformed dry weight increased linearly with  $\log_{10}$  transformed shell height for the three species, but *B. nyassanus* were heavier than either *B. globosus* or *M. tuberculata* when comparing snails of similar size.

Generally, as the shell height increased, so did CVI (Figure 1), although the slopes differed significantly ( $p < 0.001$ ), and the slope for *B. nyassanus* was not significantly different from zero. *Bulinus globosus* had the highest CVI at a given size. *Melanoides tuberculata*, however, showed a much steeper increase in CVI with shell size than the two *Bulinus* species. Within the size range covered by *B. nyassanus*, it had a higher CVI than *M. tuberculata* and only when *M. tuberculata* exceeded a shell height of about 15 mm did CVI on average exceed that of the smaller sized *B. nyassanus*. Since fish tend to crush snails against their smaller dimension (Stein et al. 1984), the shell width might be a better representation of shell size than height. When CVI was plotted against shell width (data not shown), there was no indication of a difference between *B. nyassanus* and *M. tuberculata*.

Table 1. Characteristics of the three snail species investigated.

	<i>Bulinus globosus</i> ( $n = 59$ )	<i>Bulinus nyassanus</i> ( $n = 58$ )	<i>Melanoides tuberculata</i> ( $n = 70$ )
Shell height range (mm)	5.8–16.5	4.8–11.1	8.6–28.8
Dry weight range (mg)	13.3–251.2	12.0–242.9	17.1–554.1
Organic weight range (mg)	3.5–48.0	1.2–65.7	2.3–65.2
Crush resistance range (kg)	0.234–0.489	0.85–12.014	1.90–9.669
Crush value resistance (CVI)	19.8–14.9	1.5–5.5	1.2–6.7




Figure 1. CVI of *B. nyassanus*, *B. globosus*, and *M. tuberculata* in relation to shell height.

*Bulinus globosus* has the highest CVI, which might suggest that it would be the most profitable prey for *T. placodon* followed by *B. nyassanus* and *M. tuberculata*. The habitat of *B. globosus* within the lake (sheltered areas with macrophytes or stones) only slightly overlaps with that of *T. placodon* and therefore, this snail species is not consumed in any appreciable numbers by *T. placodon* (Evers et al. 2006). There may be other predators of *B. globosus*, however, such as *Metriaclima lanisticola*, which is an oral sheller (Lundeba et al. 2007), and hence crush resistance is of little significance. The habitat of *B. globosus* may be in refugia provided by heavy vegetation, while *B. nyassanus* with its stronger shell and its behavior of descending into the top sediment is capable of choosing open sand habitats where it is more vulnerable to molluscivores. Although *B. nyassanus* has higher CVI than small *M. tuberculata* and therefore is potentially a better catch, the higher environmental abundance of *M. tuberculata* might make this species the most commonly consumed prey. The disproportionate relative abundance of *B. nyassanus* in the stomachs of *T. placodon* (Evers et al. 2006) however, would indicate preferential foraging by *T. placodon*.

Differences in shell morphology between *Bulinus* spp. and *M. tuberculata* are complicating comparisons between these species. If *T. placodon*, like some molluscivores (Stein et al. 1984), manipulates snails between pharyngeal plates to crush them against their minimal dimension, the shell size and crush resistance are not the only determinants of prey value. The globose shell of *B. nyassanus* compared to the elongated shell of *M. tuberculata* may make it easier to handle for a predator. Thus, shell shape and other morphological characteristics may function as anti-predator adaptations (Vermeij and Currey 1980; West et al. 1991; West and Cohen 1996).

Predictions about molluscivores' prey of choice are difficult to make and are influenced by other factors than the preys' crushing resistance. Factors such as spatial distribution, dispersion, position in sediment, vegetation attachment, shell morphology, and refugia activity also need consideration (Palmer 1979; Osenberg and Mittelbach 1989; Alexander and Covich 1991). The possible implications of increased density of molluscivore fishes for populations of the intermediate hosts

within the lake therefore are difficult to predict. The primary snail predator along exposed shorelines with sandy sediment is *T. placodon*, and in this habitat *B. nyassanus* is the principal host for *S. haematobium*. It is interesting that *T. placodon* seems to preferably consume *B. nyassanus* over *Melanoides* spp. (Evers et al. 2006), possibly reflecting the larger prey value of *B. nyassanus* as shown in this study.

It is possible that *T. placodon* selects indiscriminately and thus is an opportunistic snail predator. Density of *T. placodon* in shallow water (to a depth of about 5 m) at Chembe was clearly lower in 2003 than it was in 1980, and *B. nyassanus* density peaked at water depths of 1.5–6 m (Stauffer et al. 2006). Preventing beach seine-net fishing would seem to be a realistic approach for snail control in Lake Malaŵi to reduce the incidence of schistosomiasis.

### Acknowledgements

Funding was provided by the NSF/NIH joint program in Ecology of Infectious Diseases (DEB-0224958). The work was conducted in collaboration with Malaŵi Fisheries Department, Malaŵi Park and Wildlife Department and Bunda College of Agriculture, University of Malaŵi.

### References

- Alexander JE, Covich AP. 1991. Predation risk and avoidance behaviour in two freshwater snails. *Biological Bulletin*. 180(3):387–393.
- Begon M, Harper JL, Townsend CR. 1996. *Ecology*. 3rd ed. London, UK: Blackwell Science Ltd.
- Brodersen J, Chimbari MJ, Madsen H. 2003. Prosobranch mollusc species- and size-preference of *Sargochromis codringtonii* (Cichlidae) in Lake Kariba, Zimbabwe. *African Journal of Aquatic Science*. 28(2):1–4.
- Cetron MS, Chitsulo L, Sullivan JJ, Pilcher J, Wilson M, Noh J, Tsang VC, Hightower AW, Addiss DG. 1996. Schistosomiasis in Lake Malawi. *The Lancet*. 348(9037):1274–1278.
- Evers BNH, McKaye KM, Stauffer Jr JR. 2006. The schistosome intermediate host, *Bulinus nyassanus*, is a preferred food for the cichlid fish, *Trematocranus placodon*, at Cape Maclear, Lake Malaŵi. *Annals of Tropical Medicine and Parasitology*. 100(1):75–85.
- Genner MJ, Michel E, Erpenbeck D, De Voogd N, Witte F, Pointier JP. 2004. Camouflaged invasion of Lake Malawi by an Oriental gastropod. *Molecular Ecology*. 13(8):2135–2141.
- Lundeba M, Likongwe JS, Madsen H, Stauffer Jr JR. 2007. Potential of *Metriaclima lanisticola* (Teleostei: Cichlidae) for biological control of schistosome intermediate host snails. *African Zoology*. 42(1):45–49.
- Madsen H, Bloch P, Phiri H, Kristensen TK, Furu P. 2001. *Bulinus nyassanus* is an intermediate host for *Schistosoma haematobium* in Lake Malaŵi. *Annals of Tropical Medicine and Parasitology*. 95(4):353–360.
- Madsen H, Stauffer JR, Bloch P, Konings A, McKaye KR, Lilongwe JS. 2004. *Schistosomiasis haematobium* transmission in Lake Malaŵi. *African Journal of Aquatic Science*. 29(1):117–119.
- Osenberg CW, Mittelbach GG. 1989. Effects of body size on the predator-prey interaction between pumpkinseed sunfish and gastropods. *Ecological Monographs*. 59(4):405–432.
- Palmer AR. 1979. Fish predation and the evolution of gastropod shell sculpture: experimental and geographical evidence. *Evolution*. 33(2):697–713.

- Pyke GH, Pulliam HR, Charnov EL. 1977. Optimal foraging: a selective review of theory and tests. *Quarterly Review of Biology*. 52(2):137–154.
- Stauffer JR, Arnegard ME, Cetron M, Sullivan JJ, Chitsulo LA, Turner GF, Chiota S, McKaye KR. 1997. Controlling vectors and hosts of parasitic diseases using fishes. *BioScience*. 47(1):41–49.
- Stauffer Jr JR, Madsen H, McKaye H, Konings A, Bloch P, Ferreri CP, Likongwe J, Makaula P. 2006. Molluscivorous fishes – potential for biological control of schistosomiasis. *EcoHealth*. 3:22–27.
- Stein RA, Goodman CG, Marschall EA. 1984. Using time and energetic measures of cost in estimating prey value for fish predators. *Ecology*. 65(3):702–715.
- Vermeij DJ, Currey JD. 1980. Geographical variation in the strength of thaidid snail shells. *Biological Bulletin*. 158(3):383–389.
- West K, Cohen A. 1996. Shell microstructure of gastropods from Lake Tanganyika: adaptation, convergent evolution and escalation. *Evolution*. 50(2):672–681.
- West K, Cohen A, Baron M. 1991. Morphology and behavior of crabs and gastropods from Lake Tanganyika, Africa: implications for lacustrine predator-prey coevolution. *Evolution*. 45(3):589–607.
- WHO. 1993. The control of schistosomiasis. Second report of the WHO Expert Committee. Geneva: WHO Library Cataloguing-in-Publication Data. Technical support series 830.